

READ THIS INFORMATION CAREFULLY BEFORE USE. KEEP THIS LEAFLET FOR REFERENCE.

1. MARKETING AUTHORISATION HOLDER AND MANUFACTURER RESPONSIBLE FOR BATCH RELEASE

Krka, d. d., Novo mesto
Šmarješka cesta 6
8501 Novo mesto, Slovenia

2. NAME OF THE VETERINARY MEDICINAL PRODUCT

Anthelmin[®] Plus Flavour tablets for dogs

3. STATEMENT OF THE ACTIVE SUBSTANCE(S) AND OTHER INGREDIENT(S)

Each tablet contains:

Active substances:

Praziquantel	50 mg
Pyrantel embonate	144 mg
Febantel	150 mg

Round, yellow, uncoated tablets with visible darker spots and bevelled edges with cross line on one side and plain on other side.

The tablets can be divided into equal halves and equal quarters.

4. INDICATION(S)

For the treatment of mixed infestations with the following roundworms and tapeworms in adult dogs and puppies:

Nematodes:

Ascarids: *Toxocara canis*, *Toxascaris leonina* (late immature forms and mature forms)

Hookworms: *Uncinaria stenocephala*, *Ancylostoma caninum* (adults)

Cestodes:

Tapeworms: *Taenia* spp., *Dipylidium caninum*

5. CONTRAINDICATIONS

Do not use simultaneously with piperazine compounds.

Do not exceed the stated dosage when treating pregnant bitches.

Do not use in animals with a known hypersensitivity to the active substance or to any of excipients.

Not for use in dogs younger than 2 weeks of age and/or weighing less than 2 kg.

6. ADVERSE REACTIONS

In very rare cases transient loose faeces, diarrhoea and/or vomiting may occur.

If you notice any serious effects or other effects

not mentioned in this leaflet, please inform your veterinary surgeon.

7. TARGET SPECIES

Dogs (small and medium size)

8. DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF ADMINISTRATION

For oral administration.

The recommended dose rates are: 15 mg/kg bodyweight febantel, 14.4 mg/kg pyrantel and 5 mg/kg praziquantel. This is equivalent to 1 tablet per 10 kg bodyweight.

Tablets may be halved/quartered to allow accuracy of dosing.

Body weight	Anthelmin [®] Plus Flavour tablets for dogs
Puppies and Small Dogs:	
3–5 kg	½ tablet
6–10 kg	1 tablet
Medium Dogs:	
11–15 kg	1 ½ tablets
16–20 kg	2 tablets
21–25 kg	2 ½ tablets
26–30 kg	3 tablets
Large Dogs:	
31–35 kg	3 ½ tablets
36–40 kg	4 tablets

The tablet(s) can be given directly to the dog or disguised in food. No restriction of access to food is required either before or after administration of the product.

9. ADVICE ON CORRECT ADMINISTRATION

To ensure administration of a correct dose, body weight should be determined as accurately as possible.

Puppies may be wormed with this product from 2 weeks of age and every 2 weeks until 12 weeks of age. Thereafter they should be treated at 3 monthly intervals until 6 months of age.

For the control of *Toxocara*, nursing bitches should be dosed 2 weeks after giving birth and every 2 weeks until weaning.

For adult dogs, a single dose should be used.

The advice of a veterinarian should be sought regarding the need for and frequency of repeat treatment.

In the event of a heavy roundworm infestation, a

repeat dose should be given after 14 days.

10. WITHDRAWAL PERIOD

Not applicable.

11. SPECIAL STORAGE PRECAUTIONS

Keep out of the sight and reach of children.

This medicinal product does not require any special storage conditions.

Do not use this veterinary medicinal product after the expiry date which is stated on the packaging after EXP.

12. SPECIAL WARNING(S)

Special warnings for each target species:

Fleas serve as intermediate hosts for one common type of tapeworm – *Dipylidium caninum*. Tapeworm infestation is certain to re-occur unless control of intermediate hosts such as fleas, mice etc. is undertaken.

Parasite resistance to any particular class of anthelmintic may develop following frequent, repeated use of an anthelmintic of that class.

Special precautions for use in animals:

Any part-used tablets should be discarded.

Not for use in dogs younger than 2 weeks and/or weighing less than 2 kg.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

In the interests of good hygiene, persons administering the tablet directly to a dog or by adding it to the dog's food, should wash their hands afterwards.

In case of accidental ingestion, seek medical advice and show this leaflet to the physician.

Use during pregnancy, lactation or lay:

Consult a veterinary surgeon before treating pregnant animals for roundworms.

The tablets may be used during lactation.

Do not use in bitches during the first two-thirds of pregnancy.

Interaction with other medicinal products and other forms of interaction:

Do not use simultaneously with piperazine as the anthelmintic effects of pyrantel and piperazine (used in many worming products for dogs) may be antagonized.

Concurrent use with other cholinergic compounds can lead to toxicity.

Overdose (symptoms, emergency procedures,

antidotes), if necessary:

Benzimidazoles possess wide safety margin.

Pyrantel is not absorbed systematically to any extent. Praziquantel also has a wide safety margin, of up to five times the recommended dose.

13. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCT OR WASTE MATERIALS, IF ANY

Medicines should not be disposed of via wastewater. Dispose of any unused product and empty containers in accordance with guidance from your local waste regulation authority.

14. DATE OF LEAFLET REVISION - June 2018

15. OTHER INFORMATION

For any information about this veterinary medicinal product, please contact the local representative of the marketing authorisation holder.

Local representative:

Krka UK Ltd
Churchill House
London Road
Slough SL3 7FJ
Tel: 02071 646156
Email: info.uk@krka.biz

UK only

NFA-VPS To be supplied by a veterinarian, pharmacist or Suitably Qualified Person
Marketing Authorisation No.: Vm 01656/4015